

HARD PRODUCTION OF EXOTIC HYBRID MESONS

I. ANIKIN^{1,2,5}, B. PIRE², L. SZYMANOWSKI^{3,4}, O.V. TERYAEV¹ and S. WALLON⁵


¹ *Bogoliubov Laboratory of Theoretical Physics, JINR, 141980 Dubna, Russia*

² *CPHT^a, École Polytechnique, 91128 Palaiseau, France*

³ *Soltan Institute for Nuclear Studies, Warsaw, Poland*

⁴ *Université de Liège, B4000 Liège, Belgium*

⁵ *LPT^b, Université Paris-Sud, 91405-Orsay, France*



Exotic hybrid mesons H , with quantum numbers $J^{PC} = 1^{-+}$ may be copiously produced in the hard exclusive processes $\gamma^*(Q^2)\gamma \rightarrow H$ and $\gamma^*(Q^2)P(p) \rightarrow HP(p')$ because they have a leading twist distribution amplitude with a sizable coupling constant f_H , which may be estimated through QCD sum rules. The reaction rates scale in the same way as the corresponding rates for usual mesons.

1 Introduction

Within quantum chromodynamics, hadrons are described in terms of quarks, anti-quarks and gluons. The usual, well-known, mesons are supposed to contain quarks and anti-quarks as valence degrees of freedom while gluons play the role of carrier of interaction. On the other hand, QCD does not prohibit the existence of the explicit gluonic degree of freedom in the form of a vibrating flux tube, for instance. The states where the $q\bar{q}g$ and gg configurations are dominating, hybrids and glueballs, are of fundamental importance to understand the dynamics of quark confinement and the nonperturbative sector of quantum chromodynamics. We investigated¹ how hybrid mesons with $J^{PC} = 1^{-+}$, one candidate being $\pi_1(1400)$, may be studied through the so-called hard reactions. We focussed on deep exclusive γ^*P reactions which are well described in the framework of the collinear approximation where generalized parton distributions (GPDs)² and (generalized) distribution amplitudes^{3,4} describe the nonperturbative parts of a factorized

^aUnité mixte C7644 du CNRS

^bUnité mixte 8627 du CNRS

amplitude^{3,5}. We now suggest to use $\gamma^*\gamma$ reactions at e^+e^- colliders to reveal such hybrid mesons.

2 Hybrid meson distribution amplitude

Let us consider the properties of the longitudinally polarized hybrid meson distribution amplitude. The Fourier transform of the hybrid meson-to-vacuum matrix element of the bilocal vector quark operator may be written as

$$\langle H_L(p, 0) | \bar{\psi}(-z/2) \gamma_\mu \psi(z/2) | 0 \rangle = i f_H M_H e_{L\mu}^{(0)} \int_0^1 dy e^{i(\bar{y}-y)p \cdot z/2} \phi_L^H(y), \quad (1)$$

where $e_{L\mu}^{(0)} = (e^{(0)} \cdot z) / (p \cdot z) p_\mu$ and $\bar{y} = 1 - y$ and H denotes the isovector triplet of hybrid mesons; f_H denotes a dimensionful coupling constant of the hybrid meson, so that ϕ^H is dimensionless.

In (1), we imply the path-ordered gluonic exponential along the straight line connecting the initial and final points $[z_1; z_2]$ which provides the gauge invariance for bilocal operator and equals unity in a light-like (axial) gauge. For simplicity of notation we shall omit the index L from the hybrid meson distribution amplitude.

Although exotic quantum numbers like $J^{PC} = 1^{-+}$ are forbidden in the quark model, it does not prevent the leading twist correlation function from being non zero. The basis of the argument is that the non-locality of the quark correlator opens the possibility of getting such a hybrid state, because of dynamical gluonic degrees of freedom arising from the Wilson line (more details can be found in¹).

To estimate the magnitude of the distribution amplitude, one uses the estimate of the matrix element of the quark energy-momentum tensor between vacuum and exotic meson state which was explored long ago⁶. It may be related, by making use of the equations of motion, to the matrix element of quark-gluon operator and estimated with the help of the techniques of QCD sum rules. One of the solutions corresponds to a resonance with mass around 1.4 GeV and the coupling constant $f_H \approx 50$ MeV.

In summary, the hybrid light-cone distribution amplitude is a leading twist quantity which should have a vanishing first moment because of its symmetry properties. It obeys usual evolution equations³ and has an asymptotic limit

$$\Phi_{as}^H = 30y(1-y)(1-2y), \quad (2)$$

which allows to fix the value of the coupling constant f_H assuming normalization of the distribution amplitude $\Phi^H(y)$ as

$$\int_0^1 dy (1-2y) \Phi^H(y) = 1.$$

3 $\gamma^*(Q^2)\gamma \rightarrow H$

The reaction $\gamma^*(Q^2)\gamma \rightarrow H$ may be studied at intense electron-positron colliders such as CLEO, BABAR and BELLE. The production amplitude is proportional to the hybrid meson distribution amplitude $\Phi_H(z)$. It reads

$$T_{\mu\nu}(\gamma\gamma^* \rightarrow H_L) = g_{\mu\nu}^T \frac{(e_u^2 - e_d^2) f_H}{2\sqrt{2}} \int_0^1 dz \Phi^H(z) \left(\frac{1}{\bar{z}} - \frac{1}{z} \right), \quad (3)$$

with $g_{\mu\nu}^T$ projecting on the transverse coordinate subspace. It is convenient to study the cross-section ratio

$$\frac{d\sigma^H}{d\sigma^{\pi^0}} = \left| \frac{f_H \int_0^1 dz \Phi^H(z) \left(\frac{1}{z} - \frac{1}{\bar{z}}\right)}{f_\pi \int_0^1 dz \Phi^\pi(z) \left(\frac{1}{z} + \frac{1}{\bar{z}}\right)} \right|^2. \quad (4)$$

We want to stress that using asymptotic distribution amplitude for the hybrid meson (2) and for the pion $\Phi^\pi(z) = 6z\bar{z}$, the obtained ratio of cross sections is quite large, $d\sigma^H/d\sigma^{\pi^0} \approx 38\%$. If the process of reference is taken as $\gamma^*\gamma \rightarrow \eta$ then the corresponding ratio is $d\sigma^H/d\sigma^\eta \approx 46\%$.

One may have to base the hybrid identification process through one of its possible decay products. If the $\pi\eta$ decay mode is dominant as seen for $\pi_1(1400)$ (or $\pi\eta'$ for the candidate $\pi_1(1600)$), we need to introduce the concept of generalized distribution amplitude (GDA) for $\pi\eta$. This factorized description of exclusive processes in $\gamma^*\gamma$ collision has been shown⁷ to be successful for Q^2 greater than a few GeV^2 . The $\pi^0\eta$ GDA may be defined as :

$$\langle \pi^0(p_\pi)\eta(p_\eta) | \bar{\psi}_{f_2}(-z/2)\gamma^\mu[-z/2; z/2]\tau_{f_2 f_1}^3 \psi_{f_1}(-z) | 0 \rangle = p_{\pi\eta}^\mu \int_0^1 dy e^{i(\bar{y}-y)p_{\pi\eta}\cdot z/2} \Phi^{(\pi\eta)}(y, \zeta, m_{\pi\eta}^2), \quad (5)$$

where the total momentum of $\pi\eta$ pair $p_{\pi\eta} = p_\pi + p_\eta$, while $m_{\pi\eta}^2 = p_{\pi\eta}^2$. This $\pi\eta$ distribution amplitude $\Phi^{(\pi\eta)}$ describes non resonant as well as resonant contributions. The ζ -parameter is defined as :

$$\zeta = \frac{p_\pi^+}{(p_\pi + p_\eta)^+} - \frac{m_\pi^2 - m_\eta^2}{2m_{\pi\eta}^2}, \quad (6)$$

which is related to the angle θ , defined as the polar angle of the π meson in the center of mass frame of the meson pair, through:

$$2\zeta - 1 = \frac{\lambda(m_{\pi\eta}^2, m_\eta^2, m_\pi^2)}{m_{\pi\eta}^2} \cos \theta. \quad (7)$$

Since the mass region around 1400 MeV is dominated by the strong $a_2(1329) (2^{++})$ resonance, it is interesting to look for the interference of the amplitudes of hybrid and a_2 production, which is linear, rather than quadratic in the hybrid electroproduction amplitude. We model the $\pi\eta$ distribution amplitude in the following form:

$$\Phi^{(\pi\eta)}(y, \zeta, m_{\pi\eta}^2) = 30y(1-y)(2y-1) \left[B_{11}(m_{\pi\eta}^2) P_1(\cos \theta) + B_{12}(m_{\pi\eta}^2) P_2(\cos \theta) \right], \quad (8)$$

with the coefficient functions $B_{11}(m_{\pi\eta}^2)$ and $B_{12}(m_{\pi\eta}^2)$ related to corresponding Breit-Wigner amplitudes when $m_{\pi\eta}^2$ is in the vicinity of $M_{a_2}^2$, M_H^2 .

4 Deep exclusive electroproduction

The exotic hybrid meson may also be studied by means of its deep exclusive electroproduction, *i.e.*

$$\gamma_L^*(q) + N(p_1) \rightarrow H_L(p) + N(p_2) \quad (9)$$

when the baryon is scattered at small angle and the transferred momentum $-q^2 = Q^2$ is large (Bjorken regime). Within this regime, a factorization theorem is valid at the leading twist level, and a partonic subprocess part described in perturbative QCD is convoluted with universal soft

parts, which are GPDs and meson distribution amplitudes. The leading order amplitude for the process (9) is

$$\mathcal{A}_{(q)} = \frac{e\pi\alpha_s f_H C_F}{\sqrt{2}N_c Q} \left[e_u \mathcal{H}_{uu}^- - e_d \mathcal{H}_{dd}^- \right] \mathcal{V}^{(H, -)}, \quad (10)$$

where

$$\begin{aligned} \mathcal{H}_{ff}^\pm &= \int_{-1}^1 dx \left[\overline{U}(p_2) \hat{n} U(p_1) H_{ff'}(x) + \overline{U}(p_2) \frac{i\sigma_{\mu\alpha} n^\mu \Delta^\alpha}{2M} U(p_1) E_{ff'}(x) \right] \left[\frac{1}{x + \xi - i\epsilon} \pm \frac{1}{x - \xi + i\epsilon} \right], \\ \mathcal{V}^{(M, \pm)} &= \int_0^1 dy \phi^M(y) \left[\frac{1}{y} \pm \frac{1}{1-y} \right]. \end{aligned} \quad (11)$$

Here, functions H and E are standard leading twist GPD's. In (11), we include the definition of $\mathcal{H}_{ff}^+$ and $\mathcal{V}^{(M, +)}$ which are useful for the comparison with the ρ meson case. We discussed in details elsewhere¹ the resulting cross section. Let us just quote here the estimate for the ratio of hybrid and ρ meson electroproduction cross-sections:

$$\frac{d\sigma^H(Q^2, x_B, t)}{d\sigma^\rho(Q^2, x_B, t)} = \left| \frac{f_H}{f_\rho} \frac{(e_u \mathcal{H}_{uu}^- - e_d \mathcal{H}_{dd}^-) \mathcal{V}^H}{(e_u \mathcal{H}_{uu}^+ - e_d \mathcal{H}_{dd}^+) \mathcal{V}^\rho} \right|^2 \approx 13\% \quad \text{for } x_B = 0.33. \quad (12)$$

For smaller x_B , e.g. for $x_B = 0.18$, this ratio of cross sections is roughly 3%.

5 Conclusion

In conclusion, we have calculated the leading twist contribution to exotic hybrid meson with $J^{PC} = 1^{-+}$ production amplitude in the deep exclusive region. The resulting order of magnitude of the cross sections is sizeable and should be measurable at dedicated experiments at BABAR and BELLE on the one hand, and at JLab, Hermes or Compass on the other hand. We also estimated the effects of next to leading order contributions⁸. In the region of small Q^2 higher twist contributions should be carefully studied and included. We leave this study for future works.

Work of L.Sz. is supported by the Polish Grant 1 P03B 028 28. He is a Visiting Fellow of the FNRS (Belgium). I.V.A. thanks NATO for a grant.

1. I. V. Anikin *et al.*, Phys. Rev. D **71**, 034021 (2005) and Phys. Rev. D **70**, 011501 (2004).
2. D. Müller *et al.*: Fortsch. Phys. **42** (1994) 101; X. Ji: Phys. Rev. Lett. **78** (1997) 610; A. V. Radyushkin: Phys. Rev. D **56** (1997) 5524 ; for a recent review, see M. Diehl, Phys. Rept. **388** (2003) 41 and references therein.
3. G.P. Lepage and S.J. Brodsky: Phys. Lett. **B87** (1979) 359; A.V. Efremov and A.V. Radyushkin: Phys. Lett. **B94** (1980) 245.
4. M. Diehl *et al.*, Phys. Rev. Lett. **81** (1998) 1782; Phys. Rev. D **62** (2000) 073014.
5. J. C. Collins, L. Frankfurt, M. Strikman: Phys. Rev. D **56** (1997) 2982.
6. I. I. Balitsky *et al.*, Phys. Lett. B **112** (1982) 71; Z. Phys. C **33** (1986) 265.
7. I. V. Anikin *et al.*, Phys. Rev. D **69**, 014018 (2004) and arXiv:hep-ph/0506277.
8. I. V. Anikin *et al.*, Eur. Phys. J. C **42**, 163 (2005).